

Prevention & Treatment

What can I do to protect myself from getting sick?

There is no vaccine available right now to protect against 2009 H1N1 virus. However, a 2009 H1N1 vaccine is currently in production and may be ready for the public in the fall. As always, a vaccine will be available to protect against seasonal influenza. There are everyday actions that can help prevent the spread of germs that cause respiratory illnesses like influenza.

Take these everyday steps to protect your health:

- Cover your nose and mouth with a tissue when you cough or sneeze. Throw the tissue in the trash after you use it.
- Wash your hands often with soap and water. If soap and water are not available, use an alcohol-based hand rub.*
- Avoid touching your eyes, nose or mouth. Germs spread this way.
- Try to avoid close contact with sick people.
- If you are sick with flu-like illness, CDC recommends that you stay home for at least 24 hours after your fever is gone except to get medical care or for other necessities. (Your fever should be gone without the use of a fever-reducing medicine.) Keep away from others as much as possible to keep from making others sick.

Other important actions that you can take are:

- Follow public health advice regarding school closures, avoiding crowds and other social distancing measures.
- Be prepared in case you get sick and need to stay home for a week or so; a supply of over-the-counter medicines, alcohol-based hand rubs,* tissues and other related items might be useful and help avoid the need to make trips out in public while you are sick and contagious



What is the best way to keep from spreading the virus through coughing or sneezing?

If you are sick with flu-like illness, CDC recommends that you stay home for at least 24 hours after your fever is gone except to get medical care or for other necessities. (Your fever should be gone without the use of a fever-reducing medicine.)

Keep away from others as much as possible. Cover your mouth and nose with a tissue when coughing or sneezing.

Put your used tissue in the waste basket. Then, clean your hands, and do so every time you cough or sneeze.

If I have a family member at home who is sick with 2009 H1N1 flu, should I go to work?

Employees who are well but who have an ill family member at home with 2009 H1N1 flu can go to work as usual. These employees should monitor their health every day, and

take everyday precautions including washing their hands often with soap and water, especially after they cough or sneeze. Alcohol-based hand cleaners are also effective.* If they become ill, they should notify their supervisor and stay home. Employees who have an underlying medical condition or who are pregnant should call their health care provider for advice, because they might need to receive influenza antiviral drugs to prevent illness. For more information please see General Business and Workplace Guidance for the Prevention of Novel Influenza A (H1N1) Flu in Workers.

What is the best technique for washing my hands to avoid getting the flu?

Washing your hands often will help protect you from germs. Wash with soap and water or clean with alcohol-based hand cleaner*. CDC recommends that when you wash your hands -- with soap and warm water -- that you wash for 15 to 20 seconds. When soap and water are not available, alcohol-based disposable hand wipes or gel sanitizers may be used. You can find them in most supermarkets and drugstores. If using gel, rub your hands until the gel is dry. The gel doesn't need water to work; the alcohol in it kills the germs on your hands.

What should I do if I get sick?

If you live in areas where people have been identified with 2009 H1N1 flu and become ill with influenza-like symptoms, including fever, body aches, runny or stuffy nose, sore throat, nausea, or vomiting or diarrhea, you should stay home and avoid contact with other people. CDC recommends that you stay home for at least 24 hours after your fever is gone except to get medical care or for other necessities. (Your fever should be gone without the use of a fever-reducing medicine.) Stay away from others as much as possible to keep from making others sick. Staying at home means that you should not leave your home except to seek medical care. This means avoiding normal activities, including work, school, travel, shopping, social events, and public gatherings.

If you have severe illness or you are at high risk for flu complications, contact your health care provider or seek medical care. Your health care provider will determine whether flu testing or treatment is needed.

If you become ill and experience any of the following warning signs, seek emergency medical care.

In children, emergency warning signs that need urgent medical attention include:

- Fast breathing or trouble breathing
- Bluish or gray skin color
- Not drinking enough fluids
- Severe or persistent vomiting
- Not waking up or not interacting
- Being so irritable that the child does not want to be held
- Flu-like symptoms improve but then return with fever and worse cough

In adults, emergency warning signs that need urgent medical attention include:

- Difficulty breathing or shortness of breath

- Pain or pressure in the chest or abdomen
- Sudden dizziness
- Confusion
- Severe or persistent vomiting
- Flu-like symptoms improve but then return with fever and worse cough

Are there medicines to treat 2009 H1N1 infection?

Yes. CDC recommends the use of oseltamivir or zanamivir for the treatment and/or prevention of infection with 2009 H1N1 flu virus. Antiviral drugs are prescription medicines (pills, liquid or an inhaled powder) that fight against the flu by keeping flu viruses from reproducing in your body. If you get sick, antiviral drugs can make your illness milder and make you feel better faster. They may also prevent serious flu complications. During the current pandemic, the priority use for influenza antiviral drugs is to treat severe influenza illness (for example hospitalized patients) and people who are sick who have a condition that places them at high risk for serious flu-related complications.

What is CDC's recommendation regarding "swine flu parties"?

"Swine flu parties" are gatherings during which people have close contact with a person who has 2009 H1N1 flu in order to become infected with the virus. The intent of these parties is for a person to become infected with what for many people has been a mild disease, in the hope of having natural immunity 2009 H1N1 flu virus that might circulate later and cause more severe disease.

CDC does not recommend "swine flu parties" as a way to protect against 2009 H1N1 flu in the future. While the disease seen in the current 2009 H1N1 flu outbreak has been mild for many people, it has been severe and even fatal for others. There is no way to predict with certainty what the outcome will be for an individual or, equally important, for others to whom the intentionally infected person may spread the virus.

CDC recommends that people with 2009 H1N1 flu avoid contact with others as much as possible. If you are sick with flu-like illness, CDC recommends that you stay home for at least 24 hours after your fever is gone except to get medical care or for other necessities. (Your fever should be gone without the use of a fever-reducing medicine.) Stay away from others as much as possible to keep from making others sick.

Contamination & Cleaning

How long can influenza virus remain viable on objects (such as books and doorknobs)?

Studies have shown that influenza virus can survive on environmental surfaces and can infect a person for 2 to 8 hours after being deposited on the surface.

What kills influenza virus?

Influenza virus is destroyed by heat (167-212°F [75-100°C]). In addition, several chemical germicides, including chlorine, hydrogen peroxide, detergents (soap), iodophors (iodine-based



antiseptics), and alcohols are effective against human influenza viruses if used in proper concentration for a sufficient length of time. For example, wipes or gels with alcohol in them can be used to clean hands. The gels should be rubbed into hands until they are dry.

***What if soap and water are not available and alcohol-based products are not allowed in my facility?** Updated on Sept 14

If soap and water are not available and alcohol-based products are not allowed, other hand sanitizers that do not contain alcohol may be useful.

What surfaces are most likely to be sources of contamination?

Germs can be spread when a person touches something that is contaminated with germs and then touches his or her eyes, nose, or mouth. Droplets from a cough or sneeze of an infected person move through the air. Germs can be spread when a person touches respiratory droplets from another person on a surface like a desk, for example, and then touches their own eyes, mouth or nose before washing their hands.

How should waste disposal be handled to prevent the spread of influenza virus?

To prevent the spread of influenza virus, it is recommended that tissues and other disposable items used by an infected person be thrown in the trash. Additionally, persons should wash their hands with soap and water after touching used tissues and similar waste.



What household cleaning should be done to prevent the spread of influenza virus?

To prevent the spread of influenza virus it is important to keep surfaces (especially bedside tables, surfaces in the bathroom, kitchen counters and toys for children) clean by wiping them down with a household disinfectant according to directions on the product label.

How should linens, eating utensils and dishes of persons infected with influenza virus be handled?

Linens, eating utensils, and dishes belonging to those who are sick do not need to be cleaned separately, but importantly these items should not be shared without washing thoroughly first.

Linens (such as bed sheets and towels) should be washed by using household laundry soap and tumbled dry on a hot setting. Individuals should avoid "hugging" laundry prior to washing it to prevent contaminating themselves. Individuals should wash their hands with soap and water or alcohol-based hand rub immediately after handling dirty laundry.

Eating utensils should be washed either in a dishwasher or by hand with water and soap.